

**FAIRFAX COLON & RECTAL SURGERY, P.C.**  
DONALD B. COLVIN, M.D., F.A.S.C.R.S.  
PAUL E. SAVOCA, M.D., F.A.S.C.R.S.  
LYNDA S. DOUGHERTY, M.D., F.A.S.C.R.S.  
DANIEL P. OTCHY, M.D., F.A.S.C.R.S.  
LAWRENCE E. STERN, M.D., F.A.S.C.R.S.  
KIMBERLY A. MATZIE, M.D.  
COLORECTAL/ANORECTAL SURGERY, COLONOSCOPY, ANORECTAL PHYSIOLOGY

### **Movi Prep for Colonoscopy**

## **PLEASE READ THE FOLLOWING INSTRUCTIONS CAREFULLY AT LEAST TWO WEEKS BEFORE YOUR PROCEDURE:**

**(PATIENT MUST FOLLOW A CLEAR LIQUID DIET THE DAY PRIOR TO YOUR  
COLONOSCOPY. SEE PAGE 3 FOR CLEAR LIQUID DIET.)**

Prescribed medications for heart disease, high blood pressure and asthma may be taken prior to your procedure, unless otherwise instructed. All other medications, including diabetic medications, should be brought along with you to be taken after your procedure.

**DO NOT** take aspirin or aspirin-type products such as Ibuprofen, Advil, Aleve, Nuprin, Motrin, Naprosyn, etc. for at least one week prior to procedure. Please do not take herbal remedies or supplements for at least one week prior to your exam. You may take Tylenol. **DO NOT** take aspirin or aspirin-type products for 1 week after your colonoscopy, if you had a biopsy or polyp removed.

Please stop Plavix 10 days prior to your procedure, or Coumadin 4 days prior. Please consult the physician who prescribed the medication to make them aware.

**DO NOT** eat foods containing seeds for five days prior to your procedure, i.e. sesame or poppy seeds, strawberries, black pepper, corn or tomato skins. Do not eat olestra potato chips. If by chance you have eaten these products, still proceed with your preparation.

Please advise the physician if you have a prosthetic heart valve or if you have a serious heart murmur. Also advise physician if you have had any cardiac procedures in the last year.

On the day of your colonoscopy you will receive intravenous sedation. **These medications will impair your driving ability. You must arrange for someone to take you home after your procedure. The driver must remain in the facility throughout your procedure and recovery. No procedure will be performed unless your driver is present. You may not drive until the next day.**

The physician will give you written instructions with regards to eating after your colonoscopy.

This packet contains the **Movi-prep**. If you are unable to complete the prep, please call the Doctor's office and speak with a staff member, after hours, the Physician on call will contact you.

**FAIRFAX COLON & RECTAL SURGERY, P.C.**

DONALD B. COLVIN, M.D., F.A.S.C.R.S.

PAUL E. SAVOCA, M.D., F.A.S.C.R.S.

LYNDA S. DOUGHERTY, M.D., F.A.S.C.R.S.

DANIEL P. OTCHY, M.D., F.A.S.C.R.S.

LAWRENCE E. STERN, M.D., F.A.S.C.R.S.

KIMBERLY A. MATZIE, M.D.

COLORECTAL/ANORECTAL SURGERY, COLONOSCOPY, ANORECTAL PHYSIOLOGY

**Movi Prep for Colonoscopy**

**You must only have be on a clear liquid diet the day prior to your colonoscopy (see page 3 for sample diet)**

**Patient Instructions – Split – Dose Regimen**

The MoviPrep carton contains 4 pouched and a disposable container for mixing. You must complete the entire prep to ensure the most effective cleansing.

**The evening before your colonoscopy beginning at: 6pm**

<p>Step 1 Mix First Dose</p>	<ul style="list-style-type: none"><li>• Empty 1 Pouch A and 1 Pouch B into the disposable container</li><li>• Add lukewarm drinking water to the top line of the container. Mix to dissolve.</li></ul> <p><b>If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.</b></p>
<p>Step 2 Drink First Dose</p>	<ul style="list-style-type: none"><li>• The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed</li><li>• Drink 16 oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep.</li></ul>

**The morning of your colonoscopy 4 hours prior to leaving home take the last dose of your prep(this takes approximately 1 hour, then have NOTHING MORE TO DRINK.**

<p>Step 3 Mix Second Dose</p>	<ul style="list-style-type: none"><li>• Empty 1 Pouch A and 1 Pouch B into the disposable container</li><li>• Add lukewarm drinking water to the top line of the container. Mix to dissolve.</li></ul> <p><b>If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.</b></p>
<p>Step 4 Drink Second Dose</p>	<ul style="list-style-type: none"><li>• The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed</li><li>• <b>Do not drink or eat anything after completing the last does of your MoviPrep.</b></li></ul>

A colonoscopy prep causes the body to lose a significant amount of fluid and can result in sickness due to dehydration. It's important that you prepare your body by drinking extra clear liquids before the prep. Stay hydrated by drinking all required clear liquids during the prep. Replenish your system by drinking clear liquids after returning home from your colonoscopy.

If you have any questions, please call our office at 703-280-2841.

## FAIRFAX COLON & RECTAL SURGERY, P.C.

DONALD B. COLVIN, M.D., F.A.S.C.R.S.

PAUL E. SAVOCA, M.D., F.A.S.C.R.S.

LYNDA S. DOUGHERTY, M.D., F.A.S.C.R.S.

DANIEL P. OTCHY, M.D., F.A.S.C.R.S.

LAWRENCE E. STERN, M.D., F.A.S.C.R.S.

KIMBERLY A. MATZIE, M.D.

COLORECTAL/ANORECTAL SURGERY, COLONOSCOPY, ANORECTAL PHYSIOLOGY

# CLEAR LIQUID DIET

This diet provides fluids that leave little residue and are easily absorbed with minimal digestive activity. This diet is inadequate in all essential nutrients and is recommended only if clear liquids are temporarily needed. **No red or purple** liquids should be consumed!

Food Group	Foods Allowed	Foods to avoid
Milk & Beverage <i>No red or purple liquids</i>	Tea and coffee (no creamer), carbonated beverages, fruit flavored drinks	Milk creamer, milk drinks
Meats & Meat Substitutes	None	All
Vegetables	None	All
Fruit & Fruit Juices	Strained fruit juices: apple, white grape, lemonade	Fruit juices with unstrained fruit
Grains & Starches	None	All
Soups	Clear broth, consommé	All others
Desserts	Clear flavored gelatin, popsicles <i>No red or purple flavors</i>	All others
Fats	None	All
Miscellaneous	Sugar, honey, syrup, clear hard candy, salt	All others

*\*The following menu is only a suggestion\**

Breakfast	Lunch	Dinner
4 oz. White grape juice	4 oz. Apple juice	4 oz. Lemonade
6 oz. Clear broth	6 oz. Clear broth	6 oz. Clear broth
JELL-O*	JELL-O	JELL-O
Tea	Tea	Tea

## **FAIRFAX COLON & RECTAL SURGERY, P.C.**

DONALD B. COLVIN, M.D., F.A.S.C.R.S.

PAUL E. SAVOCA, M.D., F.A.S.C.R.S.

LYNDA S. DOUGHERTY, M.D., F.A.S.C.R.S.

DANIEL P. OTCHY, M.D., F.A.S.C.R.S.

LAWRENCE E. STERN, M.D., F.A.S.C.R.S.

KIMBERLY A. MATZIE, M.D.

COLORECTAL/ANORECTAL SURGERY, COLONOSCOPY, ANORECTAL PHYSIOLOGY

### **INFORMATION FOR PATIENTS PREPARING FOR COLONOSCOPY**

#### **COLONOSCOPY**

A procedure on a clean, prepared colon using a flexible scope with fiber optics to visualize the entire colon (also known as the large intestine) for colon cancer screening and possibly the last part of the small intestine. It also allows for treatment such as removal of polyps, biopsies of unusual areas or control of bleeding.

#### **LENGTH OF THE PROCEDURE**

The procedure usually lasts from 20-30 minutes but can take longer if polyps are present, if the colon is excessively long or twisted, or if excessive scar tissue exists. The extent of time required from check-in to check-out is approximately 2 hours.

#### **AWARENESS DURING PROCEDURE**

The procedure is usually done with anesthesia either moderate “conscious” sedation (“twilight”) where you may fall asleep but are easily aroused or a deeper sedation with IV anesthesia with an anesthesiologist present. Which anesthesia given depends on a variety of issues including medical history and past experience with anesthesia. You may experience some cramping or “gassiness” during the procedure or after.

#### **PREPARATION**

You should receive instructions for preparing your colon in the office when discussing the procedure with your physician. It can also be found on our web site [www.fairfaxcolorectal.com](http://www.fairfaxcolorectal.com).

#### **AFTER THE PROCEDURE**

You cannot drive home after the procedure due to the IV sedation and should not drive that day. You must make arrangements to have someone there to take you home from the hospital. Even if you want to take cab home you MUST have someone with you that will be responsible for you. This is a center policy and we will not allow you to go through the procedure without a responsible person to accompany you home. This person must be prepared to stay in the facility the duration of your procedure. Someone should remain with you until the following morning. Most people can return to work the following day. You should avoid going out of town after the procedure for several days due to delayed symptoms that can occur after a rare complication.

## **FAIRFAX COLON & RECTAL SURGERY, P.C.**

DONALD B. COLVIN, M.D., F.A.S.C.R.S.

PAUL E. SAVOCA, M.D., F.A.S.C.R.S.

LYNDA S. DOUGHERTY, M.D., F.A.S.C.R.S.

DANIEL P. OTCHY, M.D., F.A.S.C.R.S.

LAWRENCE E. STERN, M.D., F.A.S.C.R.S.

KIMBERLY A. MATZIE, M.D.

COLORECTAL/ANORECTAL SURGERY, COLONOSCOPY, ANORECTAL PHYSIOLOGY

### **COMPLICATIONS**

Complications uncommonly occur. If one does occur it will be treated appropriately. This could require hospitalization, medication, additional procedure, blood transfusion or surgery. (Center discharge instruction sheets will advise you on events to look out for once you return home.)

### **ALTERNATIVES**

Alternatives to Colonoscopy include sigmoidoscopy (looks at lower colon), Barium enema (radiology study) and 3-D virtual colonoscopy (CT scan study-not yet endorsed by the American College of Surgeons) which are typically discussed in the office with your physician. Video Capsule endoscopy is a study for the small intestine not for the colon.

### **RESULTS AND FOLLOW-UP**

Visual results will be given immediately both verbally and written. Results will be discussed with you and to whom you direct. You may not remember clearly after the anesthesia so it is helpful to be able to discuss results with family members waiting for you. Pathology results will be available after 10 days and you are usually contacted by our office. If you have not heard about the results after 10 days, then contact the office for your results. A report should be sent to your referring physician. If they do not receive one let us know and we will be happy to send them one.

After a routine colonoscopy, office follow-up is not usually necessary. If you would like a follow-up appointment or other issues are involved, then schedule a follow-up office appointment.

Your next recommended colonoscopy is usually based on family history, findings at the time of colonoscopy, pathology results or other risk factors.

### **INSURANCE**

Colonoscopies are frequently covered by insurance companies. The procedure will be pre-certified by our office if your insurance company requires pre-certification. You may still be responsible for a deductible or co-payment. Certain insurance companies have restrictions on specific anesthetic agents that may be used for your procedure. It is always safest for you to check with your insurance company to eliminate doubts and surprises.

### **ADDITIONAL INFORMATION**

[www.fascrs.org](http://www.fascrs.org)

[www.aca.org](http://www.aca.org)

[www.fairfaxcolorectal.com](http://www.fairfaxcolorectal.com)